

Freilandfarne

2018



| Gattung, Art, Sorte | Merkmale | Höhe in cm | Standort | Preis SFr. |
|---|----------------------|---------------|------------|---------------|
| <i>Adiantum pedatum</i> | Moorbeet | 50 | HS, Sch | 9.30 |
| <i>Adiantum venustum</i> | Moorbeet | 30 | HS, Sch | 9.30 |
| <i>Asplenium ceterach</i> (<i>Ceterach officinarum</i>) | wintergrün | 5 | S, HS, Sch | 9.30 |
| <i>Asplenium scolopendrium</i> (<i>Phyllitis scolopendrium</i>) | wintergrün | 40 | HS, Sch | 9.30 |
| <i>Asplenium scolopendrium</i> 'Cristata' | wintergrün | 40 | HS, Sch | 11.70 |
| <i>Asplenium scolopendrium</i> 'Undulatum' (<i>A. scol.</i> 'Crispa') | wintergrün, Moorbeet | 40 | HS, Sch | 11.70 |
| <i>Asplenium trichomanes</i> | wintergrün | 15 | HS, Sch | 7.90 |
| <i>Athyrium filix-femina</i> | | 80 | HS, Sch | 7.90 |
| <i>Athyrium niponicum</i> var. <i>pictum</i> (<i>Athyrium nip.</i> 'Metallicum') | | 50 | HS | 9.30 |
| <i>Athyrium otophorum</i> 'Ok anum' | | 40 | HS | 9.30 |
| <i>Blechnum penna-marina</i> | wintergrün | 15 | HS, Sch | 9.30 |
| <i>Blechnum spicant</i> | wintergrün | 40 | HS, Sch | 9.30 |
| <i>Ceterach officinarum</i> siehe <i>Asplenium ceterach</i> | | | | |
| <i>Cyrtomium fortunei</i> | wintergrün | 60 | HS, Sch | 11.70 |
| <i>Dryopteris affinis</i> (<i>Dryopteris borrieri</i>) | wintergrün | 80 | HS, Sch | 7.90 |
| <i>Dryopteris affinis</i> 'Cristata' (<i>Dryopteris affinis</i> 'The King') | wintergrün | 60 | HS, Sch | 11.70 |
| <i>Dryopteris carthusiana</i> (<i>D. spinulosa</i>) | | 80 | HS, Sch | 9.30 |
| <i>Dryopteris cristata</i> | Moorbeet | 50 | HS, Sch | 9.30 |
| <i>Dryopteris cycadina</i> (<i>Dryopteris atrata</i>) | wintergrün | 60 | HS, Sch | 9.30 |
| <i>Dryopteris erythrosora</i> | wintergrün | 50 | HS, Sch | 9.30 |
| <i>Dryopteris filix-mas</i> | | 80 | HS, Sch | 7.90 |
| <i>Dryopteris filix-mas</i> 'Barnesii' | | 80 | HS, Sch | 11.70 |
| <i>Dryopteris filix-mas</i> 'Linearis Polydactyla' | | 70 | HS, Sch | 11.70 |
| <i>Dryopteris goldiana</i> | | 120 | HS, Sch | 9.30 |
| <i>Dryopteris sieboldii</i> | wintergrün | 60 | HS, Sch | 11.70 |
| <i>Dryopteris wallichiana</i> | wintergrün | 80 | HS, Sch | 9.30 |
| <i>Gymnocarpium dryopteris</i> (<i>Currantia dryopteris</i>) | | 30 | HS, Sch | 7.90 |
| <i>Gymnocarpium robertianum</i> (<i>Currantia robertiana</i>) | | 40 | HS, Sch | 7.90 |
| <i>Matteuccia orientalis</i> | | 90 | HS, Sch | 19.00 |
| <i>Matteuccia struthiopteris</i> | | 100 | HS, Sch | 7.90 |
| <i>Onoclea sensibilis</i> | | 40 | HS, Sch | 7.90 |
| <i>Osmunda cinnamomea</i> | | 100 | HS, Sch | 11.70 |
| <i>Osmunda claytoniana</i> | | 100 | HS, Sch | 11.70 |
| <i>Osmunda regalis</i> | | 120 | HS, Sch | 11.70 |
| <i>Osmunda regalis</i> 'Gracilis' | | 60 | HS, Sch | 11.70 |
| <i>Phyllitis</i> siehe <i>Asplenium</i> | | | | |
| <i>Polypodium vulgare</i> | wintergrün | 30 | HS, Sch | 9.30 |
| <i>Polypodium vulgare</i> 'Bifido Multifidum' | wintergrün | 30 | HS, Sch | 11.70 |
| <i>Polystichum acrostichoides</i> | wintergrün | 50 | HS, Sch | 9.30 |
| <i>Polystichum aculeatum</i> | wintergrün | 60 | HS, Sch | 9.30 |
| <i>Polystichum braunii</i> | wintergrün | 60 | HS, Sch | 9.30 |
| <i>Polystichum munitum</i> | wintergrün | 60 | HS, Sch | 9.30 |
| <i>Polystichum polyblepharum</i> | wintergrün | 60 | HS, Sch | 9.30 |
| <i>Polystichum setiferum</i> | wintergrün | 50 | HS, Sch | 9.30 |
| <i>Polystichum setiferum</i> 'Plumoso-densum' | wintergrün | 40 | HS, Sch | 11.70 |

| Gattung, Art, Sorte | Merkmale | Höhe in cm | Standort | Preis SFr. |
|------------------------------------|-----------------|-----------------------|-----------------|-----------------------|
| Polystichum setiferum 'Proliferum' | wintergrün | 40 | HS, Sch | 9.30 |
| Polystichum tsussimense | wintergrün | 30 | HS, Sch | 9.30 |
| Pteridium aquilinum | | 180 | HS, Sch | 9.30 |
| Selaginella braunii | wintergrün | 25 | HS, Sch | 11.70 |
| Thelypteris palustris | | 50 | HS | 9.30 |
| Woodsia obtusa | | 30 | HS | 9.30 |